

WARNING

KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY



► ANIMAL HEALTH

BLOATEZE

Contains 700g/L Alcohol Ethoxylate and
70g/L Ethylene Propylene Oxide Block Co-Polymer

Alcohol ethoxylate bloat treatment

An aid in the prevention and treatment of frothy
bloat in cattle by drenching, pasture spray or
trough treatment



ACUTELY TOXIC SUBSTANCE (LOW HAZARD); SKIN IRRITANT (MEDIUM HAZARD); EYE IRRITANT (VERY HIGH HAZARD);
ECOTOXIC IN THE AQUATIC ENVIRONMENT (HIGH HAZARD); ECOTOXIC TO TERRESTRIAL INVERTEBRATES (MEDIUM
HAZARD); EPA APPROVAL CODE: HSR 002006. REGISTERED PURSUANT TO THE ACVM ACT 1997 NO. A4986.
SEE WWW.FOODSAFETY.GOV.TZ FOR REGISTRATION DETAILS.

► ANIMAL HEALTH

BLOATAZE

Alcohol Ethoxylate Bloat Treatment

An aid in the prevention and treatment of frothy bloat in cattle by drenching, pasture spray or trough treatment

By law, the user must take due care seeking expert advice if necessary, to avoid unnecessary pain or distress in the treated animal when using the product in a manner other than as directed on the label.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

BARCODE

4620-015

	Mild Bloat	Moderate Bloat	Severe Bloat
FIL Bloataze per cow every 12 hours			
	4mL	7mL	10mL
Dilution Diluted mixture per cow every 12 hours			
1:4	20mL	35mL	50mL
1:9	40mL	70mL	100mL

DIRECTIONS FOR USE

GENERAL: For all methods start treatment before the risk becomes severe, using a low dose rate to accustom the stock to treatment. Rapidly increase dose until control is achieved. When Bloat ceases, back off the dose cautiously, don't just stop. Methods are roughly in order of reliability. Good stock management with controlled feeding can also help reduce the severity of Bloat.

DRENCHING: Make a premix by adding FIL BLOATAZE into water. More dilute mixtures provide better control and less likelihood of dose loss. Start dosing before the onset of a bloat challenge at a rate of 4mL per cow every 12 hours. As the bloat challenge builds up, increase the dose to 7mL. A severe bloat challenge would require a dose of 10mL per cow every 12 hours.

The stated doses are for undiluted FIL BLOATAZE. The actual dose given to the animal depends on the dilution. See the table above for examples.

Do not exceed 25mL per dose as this may cause harm to the animal.

ONCE A DAY DRENCHING: Due to the long retention time of alcohol ethoxylate once a day drenching can be successful in moderate bloat conditions.

Good herd management and observation is important. Drench with 7 to 15mL, well diluted, at the morning milking.

PASTURE SPRAYING: Mix sufficient product, in 10 litres of water, to provide 3 to 6mL of FIL BLOATAZE for each animal. Add more water to give sufficient volume to spray a 12 hour break. It is essential to evenly cover the whole break. It may be necessary to re-spray in the case of heavy rain.

TROUGH TREATMENT: Allow the ballcock to operate normally. Pre-mix the FIL BLOATAZE dose into 10 litres of water before adding to the trough, to ensure adequate mixing. Pre-treat fresh troughs by adding 30mL of FIL BLOATAZE for each 100 litres of water. Add a further 30mL for each 10 cows after each 12 hour grazing. For small troughs and large herds the dose may need to be halved and administered after 4-6 hours. Alternate treatment may be necessary in severe bloat conditions.

TO RELIEVE A SEVERELY BLOATED COW: The cow is distressed and may not be able to swallow. SLOWLY administer 10 to 20mL of FIL BLOATAZE mixed in as much water as practical, either by drench gun or from a long necked bottle. Allow time for results.



HAZARD STATEMENTS

May be harmful if swallowed, in contact with skin, or inhaled. May be fatal if swallowed into airways. Causes severe skin burns and eye damage. Very toxic to aquatic life with long lasting effects.

PREVENTION STATEMENTS

Keep out of reach of children. Read label and safety data sheet before use. Avoid breathing vapours. Wear gloves and wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Wear eye, face and clothing protection. Avoid release to the environment.

RESPONSE STATEMENTS

If medical advice is needed, have product container or label at hand. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Immediately dilute with a glass of water. Call a POISON CENTRE or doctor if you feel unwell. IF ON SKIN (or hair): Wash with plenty of soap and water. Immediately remove all contaminated clothing.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If irritation or discomfort persists call a POISON CENTRE or



doctor. Absorb spillage to prevent material damage.

STORAGE AND DISPOSAL STATEMENTS

Store in the original sealed container in a cool, safe place out of reach of children and away from other chemicals and foodstuffs. Dispose of chemicals and empty containers safely in accordance with local regulations. Avoid contamination of any water supply with chemical or empty container. Triple rinse empty containers and add rinsings to use solutions.

WARNING

Traces of detergent-based bloat remedies may be toxic to calves especially pre-weaning. Wash mixing buckets and containers thoroughly before reusing.

24 HOUR EMERGENCY RESPONSE:

**0800 243 6225
(0800 CHEMCALL)**

